

Diphtheria, Tetanus, Pertussis	<p>Four or more doses of DTaP or DT, or any combination. If all four doses were given <i>before the fourth birthday</i>, a fifth dose is <i>required</i>. If the fourth dose was administered at least six months after the third dose, and on or after the fourth birthday, a fifth dose is not required.*</p> <p>One dose of Tdap vaccine must be administered on or after the tenth birthday. **</p>
	<p>Three or more doses of IPV. <i>The FINAL dose must be administered on or after the fourth birthday</i>, regardless of the number of previous doses.***</p>
Measles, Mumps, Rubella	<p>Two doses of MMR. The first dose must be administered on or after the first birthday. The second dose must be administered at least 28 days after the first dose.</p>
Hepatitis B	<p>Three doses of hepatitis B. The second dose must be administered at least 28 days after the first dose. The third dose must be given at least 16 weeks after the first dose and at least eight weeks after the second dose. The last dose in the series (third or fourth dose) must not be administered before age 24 weeks.</p>
(Chickenpox)	<p>Two doses of varicella vaccine must be administered prior to entry. The first dose must be administered on or after the first birthday. The second dose should be administered at least three months after the first dose; however, if the second dose is administered at least 28 days after the first dose, it is considered valid.</p>
Meningococcal	<p>One dose of meningococcal (serogroup A, C, W, and Y) vaccine must be administered on or after the tenth birthday.</p> <p>Two doses of MCV4. Second dose on or after age 16 years. If the first dose was given on or after the 16th birthday, only one dose is required. ****</p>

- Vaccine should be administered according to the most recent version of the *Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger* or the *Catch-up immunization schedule for persons aged four months-18 years who start late or who are more than one month behind*, as published by the Centers for Disease Control and Prevention's Advisory Committee on Immunization Practices. Schedules are available for print or download through [www.cdc.gov/vaccines/schedules/index.html](http://www.cdc.gov/vaccines/schedules/index.html).
- Vaccine doses administered less than or equal to four days before the minimum interval or age are valid (grace period). Doses administered greater than or equal to five days earlier than the minimum interval or age are not valid doses and should be repeated when age appropriate. If MMR and varicella are given on the same day, the doses must be separated by at least 28 days with no grace period.
- For additional information, please refer to the [Ohio Revised Code 3313.67](#) \_\_\_\_\_